## **VALLEYCARE**

## OLIVE VIEW-UCLA MEDICAL CENTER/HEALTH CENTERS ANGIOGRAPHY SUITE/CARDIOLOGY DIVISION POLICY & PROCEDURE PACEMAKER/AICD IMPLANTATION

NUMBER: 1916 VERSION: 1

SUBJECT/TITLE: GOWNING AND GLOVING

MD ORDER: Yes () No (X)

**POLICY:** All scrubbed persons must wear a sterile gown and gloves when participating in

surgery.

**PURPOSE:** To create and maintain a sterile field.

**DEPARTMENTS:** All

PROCEDURE: I. GOWNS

- A. Only sterile surgical gowns made from a material that has proven to be an effective barrier against the passage of microorganisms from non-sterile to sterile areas may be worn for surgical procedures.
- B. All persons who perform the surgical scrub in preparation for surgery must don a sterile gown.
- C. The scrub person should don a sterile gown from a sterile field other than the main instrument table.
- D. Once a surgical gown is donned, it is considered sterile in front from the level of the axilla to table level and the sleeves to two inches above the elbow.
  - 1. The back of the gown is not considered sterile.
  - 2. Stockinet gown cuffs are not considered sterile once a hand has passed through them.
- E. Gowns that become contaminated are changed.

## II. GLOVES

- A. All persons who perform the surgical scrub in preparation for surgery must also wear sterile gloves.
- B. Gloves must cover the stockinet sleeve of the gown.
- C. Scrub persons use closed gloving technique for initial donning of sterile gloves.

SUBJECT/TITLE: GOWNING AND GLOVING

Policy Number: 1916
Page Number: 2

D. Gloves that develop a hole or become contaminated are changed as soon as possible using open glove technique.

References:	
2008 AORN Standards and Recommended Practices.	
Approved by: Robin Wachsner (Chief of Cardiology)	Date: 08/04/2011
Review Date: 08/04/2014	Revision Date:
Distribution: Cardiology, Cardiology/Cath Lab	
Original Date: 08/04/2011	