# VALLEYCARE OLIVE VIEW-UCLA MEDICAL CENTER/HEALTH CENTERS INPATIENT OBSTETRICS POLICY & PROCEDURE

NUMBER: 4770 VERSION: 2

SUBJECT/TITLE: NEWBORN FEEDING PROTOCOL

MD ORDER: YES [X] NO []

**POLICY:** All newborns in the Normal Newborn Nursery will be assessed by an RN prior to

the initial feeding.

Infants who are breast-fed, bottle-fed, finger-fed, cup or spoon fed, or

gavaged will be stable before and after feedings.

**PURPOSE:** To establish criteria for assessing the stability of a newborn prior to feeding.

**DEPARTMENTS:** Medicine, Nursing 3D- PostPartum/Nursery

**DEFINITIONS: PROCEDURE:** 

## **CARE DIRECTIVES**

#### **ASSESSMENT:**

- 1. Nurses need to assess the newborn's ability to tolerate feedings.
- 2. Vital signs need to be assessed prior to the initial feeding.
- 3. Feedings will be given with the following vital signs:

Heart Rate:  $\leq 160 \text{ or } \geq 100$ 

Respirations: < 60 or > 30, or with no signs of respiratory

distress.

### **INTERVENTION:**

1. Feedings will be withheld with the following vital signs:

Heart Rate: > 160 or < 100

Respirations: > 60 or < 30, or with signs of respiratory

distress.

2. Physician will be notified.

#### **EVALUATION:**

SUBJECT/TITLE: NEWBORN FEEDING PROTOCOL

Policy Number: 47
Page Number: 2

1. If feedings are tolerated, continue as ordered.

2. If feedings are not tolerated, discontinue and inform physician.

#### **EDUCATION:**

Parents will be informed and educated in the different feeding methods, including the signs and symptoms of an infant not tolerating feedings.

# **DOCUMENTATION:**

Documentation will be charted on the # 9 (Newborn Flow) and the Progress Notes.

#### **BREASTFEEDING:**

- 1. Instruct and assist with proper breastfeeding techniques (positioning and latch-on).
- 2. To breastfeed on demand (at least 8 times per 24 hours).
- 3. Refer to Breastfeeding Policy.

#### **NON-BREASTFEEDING:**

- 1. Need a physician's order or need to obtain written consent from the mother before giving formula.
- 2. Instruct mother on proper bottle-feeding techniques.
- 3. Hand feed each infant. (No propping of bottles).
- 4. Give 20 cal. formulas every two (2) to three (3) hours.
- 5. Normal intake at each feeding:
  - a.  $1^{st}$  24 hours: 5 15 ml
  - b. 24 48 hours: up to 30 ml
  - c. 48 72 hours: up to 45 ml
  - d. >72 hours: ad lib
- 6. If difficulty is experienced with the regular Nuk nipple, use preemie nipple.

References:	
Approved by: Marci Hamilton (Clinical Nurse Director)	Date: 03/20/2013
Review Date: <b>03/20</b> /2016	Revision Date:
Distribution: Medicine, Nursing 3D- PostPartum/Nursery	
Original Date: 03/20/2013	